


b.) Amendments to the Claims

Claims 1-4 (Cancelled)

5. (Currently Amended) A method of inhibiting inflammation in an individual, comprising administering to the individual a therapeutically effective amount of an imidazolidine compound represented by the following Formula (I):



or a physiologically acceptable salt thereof, wherein:

Z is

- hydrogen,
- halogen,
- hydroxy,
- COOH,
- CONH₂,
- substituted or unsubstituted lower alkyl,
- substituted or unsubstituted haloalkyl,
- substituted or unsubstituted heteroalkyl,

substituted or unsubstituted cycloalkyl,
substituted or unsubstituted polycycloalkyl,
substituted or unsubstituted lower alkenyl,
substituted or unsubstituted cycloalkenyl,
substituted or unsubstituted polycycloalkenyl,
substituted or unsubstituted lower alkoxy,
substituted or unsubstituted lower alkanoyloxy,
substituted or unsubstituted lower alkanoyl,
substituted or unsubstituted lower alkoxycarbonyl,
substituted or unsubstituted aralkyl,
substituted or unsubstituted heteroaralkyl,
substituted or unsubstituted aryl,
substituted or unsubstituted heteroaryl, or
a substituted or unsubstituted non-aromatic heterocyclic group,
or

Z and R⁶ taken together form a bond, or

Z and R^{13a} taken together form a bond;

X¹ and X² are each, independently,

hydrogen,

-CN,

-NO₂,

-SO₂R^{15a},

-SO₂NR^{15a}R^{15b},

-C(=O)-R^{15a},

$-\text{C}(=\text{O})-\text{OR}^{15a}$, or

$-\text{C}(=\text{O})-\text{NR}^{15a}\text{R}^{15b}$, wherein

R^{15a} and R^{15b} are each, independently,

hydrogen,

substituted or unsubstituted lower alkyl,

substituted or unsubstituted cycloalkyl,

substituted or unsubstituted aryl, or

substituted or unsubstituted aralkyl;

Y is

a bond,

$-(\text{C}=\text{O})-$, or

$-(\text{CR}^{16a}\text{R}^{16b})-$, wherein

R^{16a} and R^{16b} are each, independently,

hydrogen,

substituted or unsubstituted lower alkyl,

substituted or unsubstituted cycloalkyl,

substituted or unsubstituted aryl, or

substituted or unsubstituted aralkyl;

R^1 is

substituted or unsubstituted lower alkyl,

substituted or unsubstituted cycloalkyl,

substituted or unsubstituted polycycloalkyl,

substituted or unsubstituted lower alkenyl,

substituted or unsubstituted cycloalkenyl,
substituted or unsubstituted lower alkoxy,
substituted or unsubstituted lower alkanoyloxy,
substituted or unsubstituted aralkyl,
substituted or unsubstituted heteroaralkyl,
substituted or unsubstituted aryl,
substituted or unsubstituted heteroaryl, or
a substituted or unsubstituted non-aromatic heterocyclic group;

R^{2a} , R^{2b} , R^{3a} , R^{3b} , R^{4a} , R^{4b} , R^{5a} , and R^{5b} are each, independently,

hydrogen,
substituted or unsubstituted lower alkyl,
substituted or unsubstituted cycloalkyl,
substituted or unsubstituted aryl,
substituted or unsubstituted aralkyl, or
substituted or unsubstituted heteroaralkyl;

R^6 , R^7 , R^8 , and R^9 are each, independently,

hydrogen,
hydroxy,
substituted or unsubstituted lower alkyl,
substituted or unsubstituted lower alkoxy,
substituted or unsubstituted lower alkanoyl,
substituted or unsubstituted lower alkanoyloxy,
substituted or unsubstituted lower alkoxy carbonyl,

substituted or unsubstituted aryl,
substituted or unsubstituted heteroaryl,
halogen,
-CN,
-NO₂,
-C(=O)-OR^{17a},
-NR^{17a}R^{17b}, or
-C(=O)-NR^{17a}R^{17b}, wherein
R^{17a} and R^{17b} are each, independently,

hydrogen,
substituted or unsubstituted lower alkyl,
substituted or unsubstituted cycloalkyl,
substituted or unsubstituted aryl, or
substituted or unsubstituted aralkyl,
or

R^{17a} and R^{17b} taken together with the nitrogen atom to which they are bonded form a substituted or unsubstituted heterocyclic group containing at least one nitrogen atom;

R^{10a}, R^{10b}, R^{11a}, and R^{11b} are each, independently,

hydrogen,
substituted or unsubstituted lower alkyl,
substituted or unsubstituted cycloalkyl,
substituted or unsubstituted aryl,
substituted or unsubstituted aralkyl,

substituted or unsubstituted heteroaralkyl, or
substituted or unsubstituted lower alkoxyalkyl;

R^{12a} and R^{12b} are each, independently,

hydrogen,
substituted or unsubstituted lower alkyl,
substituted or unsubstituted cycloalkyl,
substituted or unsubstituted aryl,
substituted or unsubstituted aralkyl, or
substituted or unsubstituted heteroaralkyl,

or

R^{12a} and R^{12b} taken together with the nitrogen atom to which they are bonded form a substituted or unsubstituted heterocyclic group containing at least one nitrogen atom;

R^{13a} and R^{13b} are each, independently,

hydrogen,
substituted or unsubstituted lower alkyl,
substituted or unsubstituted cycloalkyl,
substituted or unsubstituted aryl,
substituted or unsubstituted aralkyl, or
substituted or unsubstituted heteroaralkyl,

wherein when p is 2 or more, multiple R^{13a} 's are independently the same or different and multiple R^{13b} 's are independently the same or different;

m is an integer from 0 to 4;
n is an integer from 0 to 6;
p is an integer from 0 to 9; and
q is an integer from ~~0 to 5~~ 0 to 5.

6. (Original) The method according to Claim 5 wherein said inflammation is a consequence of an autoimmune disease.

7. (Original) The method according to Claim 5 wherein said inflammation is a consequence of an allergic disease or condition.

8. (Original) The method according to Claim 5 wherein said inflammation is a consequence of infection.

9. (Original) The method according to Claim 8 wherein said infection is bacterial, viral, fungal or parasitic.

Claims 10-12 (Cancelled).